



C. Juplandend Mattall I C. Mattalli For [C. ped monlosa Mass) and South western plate belonging to the first cetting the form with county gibbour the latte will boy pedurbel frit with an eversed contines.







1. Cor. 2 time 3. hots flant, ast elengets; carpely separated nearly the axis. smile booder than high 1 C'Aus tini lagelen. Fruit V Compresed, Scepty vivis 2° above I below on pedrials of its own length on usually shooter; lover the fruit very northern the first dvaniet leaves statulate Join July wood, i feld I road; Jabot I inh Ligh, lear, 1-12/1/20
Were york & New Jersey, C. I Austin touth with with with him to the Menter


He South Aurica Sparis a May again - Male; \$2. C. verna dis Fruit Carger, higher their brown obourte fise the upward, sessile for begette upward; jessile lobes of for Carinate or very name of ly over between the hours the fit almost and warely beginning; styles shorter the fit almost and warely devices of feeling leavy robulate spasulate name a potente en Agif to Any from a few chate to be placed to the Children of the contract of the children of the surface of the s Dungle & No Jenny Worthwestern ... is. C. heterophylla Punk pit swellen, as brows or browser the high, deeply emayinate, thick, alund authore, sissile on debraid Tober of fuit oblasely augled, with a narrow grown separaly this styler er log as fig, erach persother, floolig leavery restite speliste of states narrows to appeliste of the land of the land of same light before the land of same light by mantitioned to like this land of same light of same times to be a submey lover one of a submey Donnitt lineage Caver (C. Gineavi land) avend mere. from Man york & Alleron I southward autumelis L. Fruit large destants / much en grassed) vivielans Jeeply and narrowly notched (ressile or subsemile, Cober of sprice blavely any broadly wright, with a very deep al mann grows between them; styles very log reflex of deady, leaves ell linear forma broader best probable at the tip. Any - lept. 3-6 inches hogh, leave, 2-5 l. long; pit vanible there or more I Sant MISSOURI BOTANICAL


GARDEN

the simplest form of the brhagidence gwikhte L Flowers solonowious solitary or several (2003) together in Theaxies of the leaves, maked on between a pair of membanaseous bracts; male flowers consisting of a single stamen bearing a four relled Vanther opening by a single. by conflerence be come one called a soper by e striple slit, female flowers assistable sunly or pedicitied assistable swary, bearing two district) writest styles; cach all with a single ovolver of ful nullike 4 lobel and 4-celled, separatory into for father onested parts. Leeds Tolety and suspended filling each out anatropous; subryo slunder straight or sightly runved in the axis a nearly of the length of the wife albumen. Italinge very wastele Thousens Shorts smooth or covered the a mirante stellate publisseure visible only and the history rope Size kante accord, to orbour tacery, as in most vater plant. (Baun. ... annuely veg soll uniform Jane 8.1 Temestres small respitore posts Moderate or curete situationale and stillets probesseum statistics of bracks, country worthest the stillets probesseums, so compile of fail country statistics, probesseums, most compile of fail country S. 2. Aughobice amphibility personneds with clougated stems, languaged floating obvated I thear leaves fut their states of the state of permention; carpels of first country son the many of prody, almost of, whatter to the terrestrict my, on eatherely submerged soul of the way Leaden subwerged perenully with crowded 93, Subauman maiform linear leaver with but, festilight with witer


0 1 2 3 4 5 6 7 8 9 10


cm copyright reserved GARDEN






Aller 1 s Calle Las ila of entry to the transmitted C. /w/rlanden C. mierrenten C. Japany Sm. C. Alaska C. defles C. mary feats C. huttellist Constitution 1.03 min D. Anglich C. Obersang la GEORGE FINGHINN PAPERS O. Allmartin MICSSIM BOLVIICAL GARDEN C. Actionshi C. pedennelat Constant many C. Lermbottel









		L. Rech. Man	AG		α .	vulgaris. Stellaria que alsine agnisinna. Vaill.	
						Stellaria aquatica - Lob. Franienlum aquaticum Stellatum Tab.	MISSOURI
			Vernox C. Sersilis. I			C. gallioides Hop. Hoffm.	0 - 6
#	1	Callitriche agnotica. Smith.	juribus conssenta orals			intermedia is is C. heterophyllot Pursh Estimate C. aquatica BA.	9
		Coulibus filiformibus don: gatis ramoiis, foliis immeris Distantibus augusto-linearibus, Dernersis outern confertis latioribus, floribus monvicis.				minima Callitriche minima. Hop. Hoffm. 3 19 19 19 19 19 19 19 19 19 19 19 19 19	3 4 5
			De oliis superioribus protis, inferioribus, interprisambine writers, capsala acuta tetraqua pedunculata.	stel	larie i	lemissions Ron - V2III. Qua lenticula palustris anyustifolia, folio in apia dissectis. V2III. luviatelis	
			1 1 1 1 -	Call	itric	he serilis. S DC.	

Clavis of Calcibrith T. Peck 1865 with len!

MISSOUR BOTANICAL


0

9

7

2

		Tech, Manch	1865			Folis radiontibus	ovato-rotundatis	ægndimmersis.	Callitriche stell	Lata, FLop
		1021			elvingato-ovatis vel- rotinadalis					
					rotinadulis	Koliis minuti	s, ovatis, aqua ?	istoentibus. Call	itriche minimus	Hopp.
			interceris pluri- forstriverviis	Callitriche vernot - Froliis	inferioribus linea	richus		- Callil	iche intermedia	a. Schx.
		Destilibus; Folis superioribus			omnibus linea	ribus		Callity	che termifolia.	Pers.
3	Limitibus	pedmentatis	apia bidenta	Lis.	Callithriche autrum	ulis Apliis 1	inearibus	Callit	ishe conferrais	er Phuide
3		1 de tes						- Callis	triche predimen	lata.DC




Specimens of Ballitziche Ecceved brown Dr. Engle man prinon to Aug. 64. 1864. (1). Callitrich verna, I. Banks of small lakes &c.


Sphorite St. Loving. May 18#4-" (2) "6. Nerne, L. Banks of small strams. Little Rock
Ark. July 1834-"
Chicago May 1848" (3). E. Verna, 2. (4) E. Nemali, Kutz.! St. Louis Mr. May 1842
Ch. Jeyer." (3). E. Engliherdtiana, h. sp. "Memphis Zem. (6) C. Englishardtiana, n.s. "Fayetterille N.W. A.K.

June 1834-" (7) C. Reploider Mutt. 1 "Galveston Island 7. Lundheima 1843" (8)6. hammlorn, Kutz. Tweeden E. Fren Ly." (9) "C. antumnali, L. Buchi Rumin " (10)"E. auchmali il st. Lettertugt Rumin. 11"C. Stapneli, sur. Amsiffe Mar, 185-1-" 12 C. puduralosa Juno 13 C. Truvals Ledin 14 (. An gove formity glelvesto Ag. 8th. 1864 6.7 Austri

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved

MISSOURI BOTANICAL GARDEN







Calle Inche annager acronly to size of Diamet of finit Alegelmain Egelm. C. Trammondi Am 0.7. mm = 0.7 C. Tenfosa Chilis 0.7 C. Asa gragi An 0.7 - 0.7-0.8 C. Deflexa (pormoto. 9 —— C. verna Ent. 1.1 9.9-1. -0.7-0.8 Colles ang la Por 1.2 Corridentali, au 1.3 C. pudrer rulation 1. 4 12-1-3 C. marrota gue 1.4 C trumata gue 1.7 C. styndig Eur 1.-6-1.8 - 1.3-1.5 Chamelata lev 1.6-1.9 1,2 C. autumalis de 1.9 1.5




Alex.Leitch, APONICARY & CHRIST. Cor. 4th & Olive Sts., St. Louis. MISSOURI BOTANICAL CAMMIN GEORGE ENGELMANN PAPERS



Calletniche terrestri, Refin, med, sper. X1. 358 ex Dec. Vind III. 71 folis omnibus approximetis brevibus oblongis obtusis subsurrulentis, frættibus sessilibus latis brevi bus emangiaetis. (I) i pehistriby et at margines nivorum Amente bruel; præsertien i Nova Jersey proventing diffusa Lom. pl. Huite Stell, I.p.4. (: Bungslig Oursh. H. bon au. I. p 3. 1024 refer to Ane grayi on varna fredtber semiliber predien - latis refers to Austini & Ane Gragi fruitibs - brevibus, emergentis refers to destini. establissen i C. Terrestri ? For N. Mn. St. is probable : Honofthe Morth & Med Section of the U.S.



MISSOURI BOTANICAL GARDEN



Misnogn Celletinh Halorrhagi deal lallitiche 1025 A legallitriche Belgium, Lardina, Polynesia, Kerguelen 1. C. obtusangula Le Gall. 2. C. Asagraci Regelin. C. heterophylla Park C. linearis Pork, C. antemaaling my may Showing, New York, Louisiana

B. Frester, argel; brother variable

3. C. verna Lin C. vernalej Kuf C. pallous Gold. C. cophorage Lendy, C. mining

Hyp C. nadicans Phensh. Cocespeton Schold C. Terrestri, Alef (ripart) C. stellet, Hoff C. aestivaler Theat. C. intermedia Scht. C. angentifolistoff. Cerope - Asia - Chili - North Aueria to Read Mean. 4. C. hamulata Rief C. automalés Holy et surt. - greentas, Falas, Sandha Ausnia, Through lange to Stalea & Sirilia I C. pedrarulata DC Western & Southern Curope 6. C. ovvidentalig Hegelen Cerba 7. C. deflexa Abdra. C. temstui Deflo? C. pedunoulosa Jux Man ? Ació Janeiro, Alkausar, Panusylvani 8 C. stagnalis frop Germany, Spain, Ity, Joure, Alexandi Budia : C. platyrongekig bet not? Jung Man! 9. C. turfosa Best Chili 10 C. macropteryx Hegeler Men Holland B. Fructing super compressus, basi'strumosey 11 C. Drummondi Regelin New Meany = piploides Mitt. B Bendvallehrehe

12. C. antumiely Li Morther Asis & Comp [as Amen's] 13 C. Truncata Gregore Southern & Western Europe to Bili. see Mittall Ft. Akauses Fm. Philos Frances vol V. 1837. p. 140 x 141 C. pedravilosa & C. peplindy Tomy Out While and Cop. III. 135 Comangehate Mettelle: Grieber by MISSOURI BOTANICAL copyright reserved cm

GARDEN

